

CHERNOFF AND TROTTER TYPE PRODUCT FORMULAS

NEKLYUDOV A.

ABSTRACT. We consider the abstract Cauchy problem $\dot{x} = Ax$, $x(0) = x_0 \in \mathcal{D}(A)$ for linear operators A on a Banach space $\mathbf{X}$. We prove uniqueness of the (local) solution of this problem for a natural class of operators A . Moreover, we establish that the solution $x(\cdot)$ can be represented as a limit $\lim_{n \rightarrow \infty} \{F(t/n)^n\}$ in the weak operator topology, where a function $F : [0, \infty) \mapsto \mathcal{L}(\mathbf{X})$ satisfies $F'(0)y = Ay$, $y \in \mathcal{D}(A)$. As a consequence, we deduce necessary and sufficient conditions that a linear operator C is closable and its closure is a generator of C_0 -semigroup. We also obtain some criteria for the sum of two generators of C_0 -semigroups to be a generator of C_0 -semigroup such that the Trotter formula is valid.

CONTENTS

1. Introduction.	1
2. Preliminaries.	2
3. Closability of operators from class $\mathcal{Z}$.	3
4. Chernoff type formula for solution of the abstract Cauchy problem.	9
5. Criteria for closure of operator to be a generator of C_0 -semigroup.	11
References	20

1. INTRODUCTION.

Chernoff's Theorem can be formulated as follows (cf. [1], [3]).

Theorem. Let F be a map from $[0, \infty)$ to the space of all continuous linear operators $\mathcal{L}(\mathbf{X})$ in a Banach space $\mathbf{X}$; Z be a densely defined linear operator in $\mathbf{X}$ and the following conditions are satisfied:

- i) $F(0) = I$ and there exists $a \in \mathbb{R}$, $M \geq 1$ such that $\|F^m(\frac{s}{n})\| \leq M \exp(\frac{am}{n}s)$, for any $n, m \in \mathbb{N}$, $s \geq 0$.
- ii) $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$.
- iii) The closure of Z exists and it is a generator of C_0 -semigroup.

Then $F(t/n)^n$ converges to $\exp(t\bar{C})$ as $n \rightarrow \infty$ in the strong operator topology uniformly with respect to $t \in [0, T]$ for each $T > 0$.

One of the applications of Chernoff's Theorem is the theory of the Feynman integrals (survey of this theory can be found in the book [8]). In the paper [5] Chernoff's Theorem

Date: March 9, 2008.

2000 Mathematics Subject Classification. Primary 34G10, 47D03, 47D60; Secondary 47D06, 47D08.

Key words and phrases. Chernoff product formulas, Trotter product formulas, C_0 -semigroups.

has been used to prove the representation of the solution of the Schrödinger equation as the Feynman integral. Similar approach also has been used in [7], [6].

The main problem of application of Chernoff's Theorem is to check condition (iii). In this work we study the class $\mathcal{Z}$ of all densely defined operators Z that satisfy only conditions (i)–(ii) of Chernoff's Theorem. We can notice that condition (iii) implies the existence of a function F satisfying conditions (i)–(ii) for a given Z . Indeed, if condition (iii) is valid, then it is enough to put $F(t) = \exp(t\bar{Z})$, $t \in [0, \infty)$. Thus, the natural problem is to find conditions that a linear operator $Z \in \mathcal{Z}$ is closable and its closure is a generator of C_0 -semigroup. It turns out that conditions (i)–(ii) are sufficient for closability of an operator Z (see Theorem 3.1). Furthermore, if there exists (local) solution of the abstract Cauchy problem $\dot{x} = \bar{Z}x$, $x(0) = x_0 \in \mathcal{D}(\bar{Z})$, then this solution is unique and can be represented as a limit $\lim_{n \rightarrow \infty} \{F(t/n)^n x_0\}$ in the weak topology $\sigma(X, X^*)$ (see Theorem 4.1). Applying these results, we prove necessary and sufficient conditions that a linear operator A is closable and its closure is a generator of C_0 -semigroup (see Theorems 5.1–5.4, Corollaries 5.3–5.4). As a consequence, we deduce criteria for the sum of two generators of C_0 -semigroups to be a generator of C_0 -semigroup such that the Trotter formula is valid (see Corollaries 5.1–5.2, 5.5).

2. PRELIMINARIES.

For any normed space $(\mathbf{E}, \|\cdot\|_{\mathbf{E}})$ (on the field $\mathbb{T} \in \{\mathbb{R}, \mathbb{C}\}$) let $\mathcal{L}(\mathbf{E})$ be the space of all bounded linear operators in $\mathbf{E}$ with the topology of pointwise convergence (strong operator topology), I be the identity operator in $\mathbf{E}$. For any linear operator A in $\mathbf{E}$ let $\mathcal{D}(A)$ be the domain of A . Linear subspaces of $\mathbf{E}$ are considered as the normed spaces with the natural norms (i. e. the norms inherited from the normed space $\mathbf{E}$). The dual space $\mathbf{E}^*$ is the Banach space of all linear continuous functionals on $\mathbf{E}$ with the norm $\|\cdot\|_{\mathbf{E}^*}$ defined by $\|f\|_{\mathbf{E}^*} = \sup_{\|x\|_{\mathbf{E}} \leq 1} f(x)$, $x \in \mathbf{E}$, for each $f \in \mathbf{E}^*$. Further, we will omit the index of the space in the norm $\|\cdot\|_{\mathbf{E}}$. For any function $S : [0, \infty) \mapsto \mathcal{L}(\mathbf{E})$ we denote by S^* the function from $[0, \infty)$ to $\mathcal{L}(\mathbf{E})$ such that $S^*(s) = (S(s))^*$ for each $s \geq 0$.

Definition 2.1. We call the (strong) derivative at the point 0 of a function $S : [0, \infty) \mapsto \mathcal{L}(\mathbf{E})$ the linear map $S'(0) : \mathcal{D}(S'(0)) \mapsto \mathbf{E}$ defined by $S'(0)\psi = \lim_{h \rightarrow 0} h^{-1}(S(h)\psi - S(0)\psi)$, $\psi \in \mathcal{D}(S'(0))$, where $\mathcal{D}(S'(0))$ is the space of all $\psi \in \mathbf{E}$ such that the limit exists.

Definition 2.2. The set $\mathcal{F}_{M,a}^{\mathbf{E}}$, $M \geq 1$, $a \in \mathbb{R}$, is the set of all functions $F : [0, \infty) \mapsto \mathcal{L}(\mathbf{E})$ that satisfy the following conditions:

- (i) $F(0) = I$ and $\|F^n(\frac{s}{n})\| \leq M \exp(\frac{am}{n}s)$ for all $n, m \in \mathbb{N}$, $s \geq 0$.
- (ii) $\mathcal{D}(F'(0))$ is dense in $\mathbf{E}$.

Definition 2.3. The set $\mathcal{F}_{\mathbf{E}}$ is the set of all functions $F : [0, \infty) \mapsto \mathcal{L}(\mathbf{E})$ for which there exists $M \geq 1$, $a \in \mathbb{R}$ such that $F \in \mathcal{F}_{M,a}^{\mathbf{E}}$.

Remark 2.1. Condition (i) of Definition 2.2 can be replaced by equivalent condition (i^*) :

- (i^*) $F(0) = I$ and $\|F^n(\frac{s}{n})\| \leq M \exp(as)$ for all $n \in \mathbb{N}$, $s \geq 0$.

Definition 2.4. The set $\mathcal{Z}$ is the set of all densely defined linear operators in $\mathbf{E}$ for which there exists $F \in \mathcal{F}_{\mathbf{E}}$ such that $\mathcal{D}(Z) \subset \mathcal{D}(F'(0))$ and $Zf = F'(0)f$ for each $f \in \mathcal{D}(Z)$.

Definition 2.5. The function $T : [0, \infty) \rightarrow \mathcal{L}(\mathbf{E})$ is called C_0 -semigroup if the following conditions are satisfied:

- 1) $T(0) = I$, $T(l + m) = T(l)T(m)$ for each $l, m \in [0, \infty)$.
- 2) The function T is continuous.
- 3) There exists $M > 1$, $a \in \mathbb{R}$ such that $\|T(s)\| \leq M \exp(as)$ for each $s \geq 0$.

Definition 2.6. The linear operator Z is called the generator of C_0 -semigroup T if Z is the (strong) derivative at the point 0 of the function T .

It is a well-known fact that there exists one-to-one correspondence between C_0 -semigroups and its generators. The following results can be found in [2]:

Statement 2.1. Let $\mathbf{E}$ be a Banach space. Assume that the function $T : [0, \infty) \rightarrow \mathcal{L}(\mathbf{E})$ is C_0 -semigroup. Then $\mathcal{D}(T^{*'}(0))$ is $*$ -dense in $\mathbf{E}^*$, where $T^{*'}(0)$ is the (strong) derivative at the point 0 of the function T^* .

Statement 2.2. Let $\mathbf{E}$ be a reflexive Banach space. Assume that the function $T : [0, \infty) \rightarrow \mathcal{L}(\mathbf{E})$ is C_0 -semigroup. Then the function T^* is also C_0 -semigroup and its generator $T^{*'}(0) = (T'(0))^*$.

In the paper we assume that $\mathbf{X}$ is a Banach space and $\mathcal{B}$ is a reflexive separable Banach space or a Hilbert space.

3. CLOSABILITY OF OPERATORS FROM CLASS $\mathcal{Z}$.

Proposition 3.1. Assume that $F \in \mathcal{F}_{\mathbf{X}}$, $t > 0$ and $\{n_k\}_{k=1}^{\infty}$ is an increasing sequence of natural numbers. Then for any separable closed linear spaces $\Phi \subset \mathbf{X}$, $\Psi \subset \mathbf{X}^*$ there exists a subsequence $\{g_k\}_{k=1}^{\infty}$ of the sequence $\{n_k\}_{k=1}^{\infty}$ such that there exists

$$(3.1) \quad \lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k s]} g, \phi)$$

for any $g \in \Phi$, $\phi \in \Psi$ and $s > 0$. Furthermore, if such subsequence $\{g_k\}_{k=1}^{\infty}$ is chosen, then the family of the functions $T_s : \Phi \times \Psi \mapsto \mathbb{T}$, $s \geq 0$, defined by

$$(3.2) \quad T_s(g, \phi) = \lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k \frac{s}{t}]} g, \phi),$$

satisfies the following conditions:

- a) $T_s(g, \phi)$ is linear w.r.t. g and ϕ and the following inequality is satisfied

$$\|T_s(g, \phi)\| \leq M \exp(as) \|g\| \|\phi\|, \quad s \geq 0.$$

- b) $T_s(g, \phi)$ is continuous w.r.t. s for any $g \in \Phi$, $\phi \in \Psi$ and the continuity is uniform w.r.t. $\phi \in \Upsilon$, where Υ is a bounded subset of Ψ .

- c) If $f \in \Phi \cap \mathcal{D}(F'(0))$ and $F'(0)f \in \Phi$ then there exists

$$(T_s(f, \phi))'_s = T_s(F'(0)f, \phi)$$

for each $s \geq 0$, $\phi \in \Psi$ uniformly w.r.t. $\phi \in \Upsilon$, where Υ is a bounded subset of Ψ .

- d) Assume $\phi \in \mathcal{D}(F^{*'}(0))$ and $F^{*'}(0)\phi \in \Psi$. Then

$$T_m(f, \phi) - T_l(f, \phi) = \int_l^m T_s(f, F^{*'}(0)\phi) ds, \quad m, l \geq 0,$$

for any $f \in \Phi$.

e) If $\phi \in \mathcal{D}(F^{*'}(0))$ and $F^{*'}(0)\phi \in \Psi$, then

$$(T_s(f, \phi))'_s = T_s(f, F^{*'}(0)\phi), \quad s \geq 0,$$

for any $f \in \Phi$.

Now we will formulate some auxiliary lemmas.

Lemma 3.1. Assume that $F \in \mathcal{F}_{M,a}^{\mathbf{X}}$, $M \geq 1$, and $a = 0$. Then there exists $\lim_{i/k \rightarrow 0} (F^i(t/k) - I)g = 0$, $i, k \in \mathbb{N}$, for any $g \in \mathbf{X}$.

Remark 3.1. The limit in Lemma 3.1 is considered as a limit w.r.t. the filter base consisting of the sets of the form $\{(i, k) \mid |\frac{i}{k}| < \epsilon, i, k \in \mathbb{N}\}$, $\epsilon > 0$.

Proof of Lemma 3.1. If $g \in \mathcal{D}(F'(0))$, then the following chain of the inequalities is satisfied:

$$\begin{aligned} \|(F^i(t/k) - I)g\| &= i \frac{t}{k} \left\| \left(\frac{F(\frac{t}{k})^{i-1} + \dots + I}{i} \right) \frac{F(\frac{t}{k})g - g}{\frac{t}{k}} \right\| \\ &\leq i \frac{t}{k} M \left\| \frac{F(\frac{t}{k})g - g}{\frac{t}{k}} \right\| \leq i \frac{t}{k} 2M \|F'(0)g\|, \end{aligned}$$

where $k > k_0(g)$. Therefore, we have

$$\lim_{i/k \rightarrow 0} (F^i(t/k) - I)g = 0.$$

If $g \notin \mathcal{D}(F'(0))$, then for any $\varepsilon > 0$ there exists g' such that $\|g - g'\| < \varepsilon/2M$ and

$$\begin{aligned} \|\limsup_{i/k \rightarrow 0} (F^i(t/k) - I)g\| &\leq \|\limsup_{i/k \rightarrow 0} (F^i(t/k) - I)(g - g')\| \\ &+ \|\limsup_{i/k \rightarrow 0} (F^i(t/k) - I)g'\| \leq \varepsilon \end{aligned}$$

Since ε is arbitrary the Lemma is proved. $\square$

We can immediately deduce the following Lemma from Lemma 3.1.

Lemma 3.2. Assume that $F \in \mathcal{F}_{M,a}^{\mathbf{X}}$, $M \geq 1$, and $a = 0$. Then for any $g \in \mathbf{X}$ there exists $\lim_{|i-l/k| \rightarrow 0} (F^i(t/k) - F^l(t/k))g = 0$, $i, l, k \in \mathbb{N}$.

Remark 3.2. The limit in Lemma 3.2 is considered as a limit w.r.t. the filter base consisting of the sets of the form $\{(i, l, k) \mid 0 < |\frac{i-l}{k}| < \epsilon, i, k, l \in \mathbb{N}\}$, $\epsilon > 0$. In the same sense we will consider the limit in Lemma 3.3.

Lemma 3.3. Assume that $F \in \mathcal{F}_{M,a}^{\mathbf{X}}$, $M \geq 1$, and $a = 0$. Then for any $g \in \mathcal{D}(F'(0))$ there exists

$$\lim_{|i-l/k| \rightarrow 0} \frac{(F^i(t/k) - F^l(t/k))g}{(i-l)t/k} - (F^{\min(i,l)}(t/k))F'(0)g = 0,$$

$i, l, k \in \mathbb{N}$.

Proof of Lemma 3.3. We have the following chain of the inequalities

$$\begin{aligned}
& \limsup_{|i-l/k| \rightarrow 0} \left\| \frac{(F^i(\frac{t}{k}) - F^l(\frac{t}{k}))g}{(i-l)t/k} - (F^{\min(i,l)}(\frac{t}{k}))F'(0)g \right\| \\
& \leq M \limsup_{|i-l/k| \rightarrow 0} \left\| \frac{(F^{|i-l|}(\frac{t}{k}) - I)g}{|i-l|\frac{t}{k}} - F'(0)g \right\| \leq M \limsup_{i/k \rightarrow 0} \left\| \frac{(F^i(\frac{t}{k}) - I)g}{i\frac{t}{k}} - F'(0)g \right\| \\
& \leq M \limsup_{i/k \rightarrow 0} \left\| \left(\frac{(F(\frac{t}{k})^{i-1} + \dots + I)}{i} \right) \left(\frac{(F(\frac{t}{k})g - g)}{\frac{t}{k}} - F'(0)g \right) \right\| \\
& + M \limsup_{i/k \rightarrow 0} \left\| \left(\frac{(F(\frac{t}{k})^{i-1} + \dots + I)}{i} \right) F'(0)g - F'(0)g \right\| \\
& \leq \limsup_{i/k \rightarrow 0} M^2 \left\| \left(\frac{(F(\frac{t}{k})g - g)}{\frac{t}{k}} - F'(0)g \right) \right\| \\
& + M \limsup_{i/k \rightarrow 0} \left\| \left(\frac{(F(\frac{t}{k})^{i-1} - I) + \dots + (I - I)}{i} \right) F'(0)g \right\| = 0,
\end{aligned}$$

where the last limit is equal to 0 by Lemma 1. $\square$

Proof of Proposition 3.1. We consider only the case $a = 0$. The case $a \neq 0$ can be easily reduced to the case $a = 0$ by examining $F(s) \exp(-as)$ instead of $F(s)$.

We can choose a subsequence $\{g_k\}_{k=1}^\infty$ of the sequence $\{n_k\}_{k=1}^\infty$ such that for any non-negative rational r there exists

$$\lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k r]} g, \phi), \quad g \in \mathbf{C}, \quad \phi \in \mathbf{D},$$

where $\mathbf{C}$ and $\mathbf{D}$ are dense countable subsets of Φ and Ψ respectively. Indeed, it is possible to find such subsequence because

$$\|\{F(t/n_k)\}^{[n_k r]}\| \leq M$$

and we have a countable number of conditions on the sequence $\{g_k\}_{k=1}^\infty$. Hence, it follows from the density of $\mathbf{C}$ in Φ (and the density of $\mathbf{D}$ in Ψ) that there exists

$$\lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k r]} g, \phi)$$

for any $g \in \Phi$, $\phi \in \Psi$. Let us show that the last limit exists for any real number $r \geq 0$. Fix $r > 0$, $z \in \Psi$ and $\epsilon > 0$. Put $\epsilon' = \epsilon/3\|z\|$. We can choose $\epsilon''(g, \epsilon')$ such that

$$\|(F^i(t/k) - F^l(t/k))g\| < \epsilon',$$

when $|(i-l)/k| < \epsilon''$, $i, l, k \in \mathbb{N}$. Indeed, the existence of ϵ'' follows from Lemma 3.2. Choose also a positive $s \in \mathbb{Q}$ such that $|r - s| < \epsilon''/2$. There exists $k_0 \in \mathbb{N}$ such that $2/g_{k_0} < \epsilon''/2$. Now it follows from the existence of the limit

$$\lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k s]} g, z)$$

that we can find $n_0 > k_0$ such that for any $l, m > n_0$, $l, m \in \mathbb{N}$, we have the following inequality

$$|(\{F(t/g_l)\}^{[g_l s]} - \{F(t/g_m)\}^{[g_m s]})g, z| < \epsilon/3.$$

Therefore, we get the following chain of the inequalities

$$\begin{aligned}
& |(((\{F(t/g_l)\}^{[g_l r]} - \{F(t/g_m)\}^{[g_m r]})g, z)| \leq |(((\{F(t/g_l)\}^{[g_l r]} \\
& - \{F(t/g_l)\}^{[g_l s]})g, z)| + |(((\{F(t/g_l)\}^{[g_l s]} - \{F(t/g_m)\}^{[g_m s]})g, z)| \\
& + |(((\{F(t/g_m)\}^{[g_m s]} - \{F(t/g_m)\}^{[g_m r]})g, z)| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 \leq \epsilon,
\end{aligned}$$

where we have used the fact that

$$|[g_n r] - [g_n s]|/g_n \leq |g_n r - g_n s| + 2/g_n \leq \epsilon'', \quad n > k_0.$$

As a consequence, we get that there exists

$$\lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k s]}g, \phi), \quad s \geq 0, \quad g \in \Phi, \quad \phi \in \Psi.$$

Thus, the first part of Proposition 3.1 is proved.

Now let us consider parts (a) – (e) of Proposition 3.1.

- (a) Part (a) immediately follows from the definition of T_s .
(b) Fix $g \in \Phi$. It follows from Lemma 3.2 that for any $\epsilon > 0$ there exist $h_0 > 0$, $k_0 \in \mathbb{N}$ such that

$$|(\{F(t/g_k)\}^{[g_k \frac{s+h}{t}]}g, \phi) - (\{F(t/g_k)\}^{[g_k \frac{s}{t}]}g, \phi)| < \epsilon \|\phi\|,$$

for any $|h| < h_0$, $h \geq -s$, $k > k_0$, $k \in \mathbb{N}$. Letting $k \rightarrow \infty$, we get the inequality

$$|T_{s+h}(g, \phi) - T_s(g, \phi)| \leq \epsilon \|\phi\|.$$

From the arbitrariness of ϵ we easily infer part (b) of Proposition 3.1.

- (c) Assume that $f \in \Phi \cap \mathcal{D}(F'(0))$, $F'(0)f \in \Phi$. It follows from Lemma 3.3 that for any $\epsilon > 0$ there exist $h_0 > 0$, $k_0 \in \mathbb{N}$ such that

$$\begin{aligned}
& |(F^{[g_k \frac{s}{t}]}(t/g_k)f, \phi) - (F^{[g_k \frac{s+h}{t}]}(t/g_k)f, \phi) - t/g_k([g_k \frac{s}{t}] - [g_k \frac{s+h}{t}]) \\
& \times (F^{\min([g_k \frac{s}{t}], [g_k \frac{s+h}{t}])}(t/g_k)F'(0)f, \phi)| < \epsilon \|\phi\| |[g_k \frac{s}{t}] - [g_k \frac{s+h}{t}]| \frac{t}{g_k}
\end{aligned}$$

for any $|h| < h_0$, $h \geq -s$, $k > k_0$, $k \in \mathbb{N}$. Letting $k \rightarrow \infty$, we get the inequality

$$\|T_s(f, \phi) - T_{s+h}(f, \phi) - hT_{\min(s, s+h)}(F'(0)f, \phi)\| \leq \epsilon |h| \|\phi\|.$$

From the arbitrariness of ϵ we deduce part (c) of Proposition 3.1.

- (d) Assume that $\phi \in \mathcal{D}(F^{*'}(0))$, $F^{*'}(0)\phi \in \Psi$. Fix $\epsilon > 0$ and $m > l$. Choose r_0, j_0 such that

$$\|F(t/g_k)^{[g_k \frac{g}{t}]}f - F(t/g_k)^{[g_k \frac{h}{t}]}f\| < \epsilon, \quad k \geq j_0, \quad |g - h| < r_0, \quad 0 \leq g, h < m.$$

The existence of such r_0, j_0 follows from Lemma 3.2. Hence, we have

$$(3.3) \quad |T_h(f, F^{*'}(0)\phi) - T_g(f, F^{*'}(0)\phi)| \leq \epsilon,$$

where $|g - h| < r_0$, $0 \leq g, h < m$. Therefore, we get

$$\begin{aligned}
(3.4) \quad & \left| \sum_{v=0}^{v_0-1} T_{l+v(m-l)/v_0}(f, F^{*'}(0)\phi)((m-l)/v_0) \right. \\
& \left. - \int_l^m T_s(f, F^{*'}(0)\phi) ds \right| \leq \epsilon |m - l|,
\end{aligned}$$

$v_0 = [(m-l)/r_0] + 1$. Choose $j_1 > j_0$ such that

$$|T_{l+v(m-l)/v_0}(f, F^{*'}(0)\phi) - (F(t/g_k)^{[g_k(l+v(m-l)/v_0)/t]} f, F^{*'}(0)\phi)| < \epsilon$$

for any $v \in \{0, 1, \dots, v_0\}$, $k > j_1$. Thus,

$$(3.5) \quad \left| \sum_{v=0}^{v_0-1} T_{l+v(m-l)/v_0}(f, F^{*'}(0)\phi) \right|$$

$$(3.6) \quad - ((m-l)/v_0) \sum_{v=0}^{v_0-1} (F(t/g_k)^{[g_k(l+v(m-l)/v_0)/t]} f, F^{*'}(0)\phi) < \epsilon |m-l|.$$

Now we notice that

$$(3.7) \quad T_m(f, \phi) - T_l(f, \phi) = \lim_{k \rightarrow \infty} (F(\frac{t}{g_k})^{[g_k \frac{m}{t}]} f - F(\frac{t}{g_k})^{[g_k \frac{l}{t}]} f, \phi)$$

$$(3.8) \quad = \lim_{k \rightarrow \infty} ((F(\frac{t}{g_k}) - I)(F(\frac{t}{g_k})^{[g_k \frac{m}{t}] - 1} + \dots + F(\frac{t}{g_k})^{[g_k \frac{l}{t}]}) f, \phi)$$

$$(3.9) \quad = \lim_{k \rightarrow \infty} (\frac{t}{g_k} (F(\frac{t}{g_k})^{[g_k \frac{m}{t}] - 1} + \dots + F(\frac{t}{g_k})^{[g_k \frac{l}{t}]}) f, \frac{t}{g_k} (F^*(\frac{t}{g_k}) - I)\phi)$$

$$(3.10) \quad = \lim_{k \rightarrow \infty} (\frac{t}{g_k} (F(\frac{t}{g_k})^{[g_k \frac{m}{t}] - 1} + \dots + F(\frac{t}{g_k})^{[g_k \frac{l}{t}]}) f, F^{*'}(0)\phi)$$

$$(3.11) \quad = \lim_{k \rightarrow \infty} \frac{t}{g_k} \sum_{s=0}^{d_k} (F(\frac{t}{g_k})^{[g_k(l+s(\frac{t}{g_k}))/t]} f, F^{*'}(0)\phi)$$

$$(3.12) \quad = \lim_{k \rightarrow \infty} \sum_{v=0}^{v_0-1} \sum_{s \in \mathbf{B}_{k,v,v_0}} \frac{t}{g_k} (F(\frac{t}{g_k})^{[g_k(l+s(\frac{t}{g_k}))/t]} f, F^{*'}(0)\phi),$$

where $d_k = [(m-l)g_k/t]$ and

$$\mathbf{B}_{k,v,v_0} = \{s \in \mathbb{N} \mid v(m-l)/v_0 \leq st/g_k < (v+1)(m-l)/v_0\}.$$

So, by inequality (3.3) and equalities (3.7) — (3.12), we get

$$(3.13) \quad |T_l(f, \phi) + \sum_{v=0}^{v_0-1} (F(\frac{t}{g_k})^{[g_k(l+v\frac{m-l}{v_0})/t]} f, F^{*'}(0)\phi) (\frac{m-l}{v_0}) - T_m(f, \phi)|$$

$$(3.14) \quad = | \lim_{k \rightarrow \infty} \sum_{v=0}^{v_0-1} \sum_{s \in \mathbf{B}_{k,v,v_0}} \frac{t}{g_k} (F(\frac{t}{g_k})^{[g_k(l+s(\frac{t}{g_k}))/t]} f, F^{*'}(0)\phi)$$

$$(3.15) \quad - (\frac{m-l}{v_0}) \sum_{v=0}^{v_0-1} (F(\frac{t}{g_k})^{[g_k(l+v\frac{m-l}{v_0})/t]} f, F^{*'}(0)\phi) |$$

$$(3.16) \quad \leq \sum_{v=0}^{v_0-1} | \lim_{k \rightarrow \infty} \sum_{s \in \mathbf{B}_{k,v,v_0}} \frac{t}{g_k} (F(\frac{t}{g_k})^{[g_k(l+s(\frac{t}{g_k}))/t]} f, F^{*'}(0)\phi)$$

$$(3.17) \quad - \frac{m-l}{v_0} (F(\frac{t}{g_k})^{[g_k(l+v\frac{m-l}{v_0})/t]} f, F^{*'}(0)\phi) | < \epsilon v_0 \frac{|m-l|}{v_0}.$$

Combining inequalities (3.4), (3.5) — (3.6) and (3.13) — (3.17), we get

$$|T_m(f, \phi) - T_l(f, \phi) - \int_l^m T_s(f, F^{*'}(0)\phi) ds| < 3\epsilon|m - l|$$

From the arbitrariness of ϵ we deduce part (d) of Proposition 3.1.

(e) Part (e) is a direct consequence of parts (b) and (d). □

Proposition 3.2. Assume that $F \in \mathcal{F}_X$. Then the operator $F'(0)$ is closable.

Proof of Proposition 3.2. It is enough to show that the existence of the limits

$$\lim_{n \rightarrow \infty} f_n = 0, \quad \lim_{n \rightarrow \infty} F'(0)f_n = h, \quad h \in X,$$

for any sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \mathcal{D}(F'(0))$, imply that $h = 0$. Indeed, it immediately follows from Lemma 1.8, p. 7 in [2]. We will argue by contradiction. Assume that $h \neq 0$. Then there exists $\phi \in X^*$ such that $(h, \phi) \neq 0$. Let $\mathbf{B}$ be the closure of the linear span of the elements $f_i, F'(0)f_i$, $i \in \mathbb{N}$. Then, according to Proposition 3.1, there exists an increasing sequence $\{g_k\}_{k=1}^\infty$, $g_k \in \mathbb{N}$, such that we can define the family of the functions $T_s : \Phi \times \Psi \mapsto \mathbb{T}$, $s \geq 0$, by formula (3.2) with $\Phi = \mathbf{B}$ and $\Psi = \{\phi\}$. Hence, we have

$$(T_s(f_i, \phi))'_s = T_s(F'(0)f_i, \phi)$$

for any $i \in \mathbb{N}$. Therefore, we get

$$(3.18) \quad T_s(f_i, \phi) - (f_i, \phi) = \int_0^s T_t(F'(0)f_i, \phi) dt, \quad i \in \mathbb{N}.$$

Letting $i \rightarrow \infty$ in expression (3.18), we get the equality

$$T_s(0, \phi) - (0, \phi) = \int_0^s T_r(h, \phi) dr = 0, \quad s \geq 0.$$

Indeed, it immediately follows from (3.18) and the uniform continuity of the family $\{T_l(\cdot, \phi) | l \in [0, s]\}$. Hence, we get

$$(3.19) \quad \left(\int_0^s T_t(h, \phi) dt \right)'_s = T_s(h, \phi) = 0, \quad s \geq 0.$$

Put $s = 0$ in (3.19). Consequently, we have

$$T_s(h, \phi) = (h, \phi) = 0.$$

Thus, we get contradiction with the assumption $h \neq 0$. □

As a consequence of Proposition 3.2, we get

Theorem 3.1. Let $F \in \mathcal{F}_X$. Assume that a linear operator Z has the domain $\mathcal{D}(Z) \subset \mathcal{D}(F'(0))$ and

$$Zf = F'(0)f,$$

for any $f \in \mathcal{D}(Z)$. Then Z is closable.

4. CHERNOFF TYPE FORMULA FOR SOLUTION OF THE ABSTRACT CAUCHY PROBLEM.

Lemma 4.1. Suppose that the assumptions of Proposition 3.1 are satisfied. Assume also that $f_i \in \mathcal{D}(F'(0)) \cap \Phi$, $F'(0)f_i \in \Phi$ for any $i \in \mathbb{N}$. Furthermore, suppose that

$$\lim_{i \rightarrow \infty} f_i = h, \quad \lim_{i \rightarrow \infty} Zf_i = g.$$

Then

$$(T_s(h, \phi))'_s = T_s(g, \phi), \quad s \geq 0, \quad \phi \in \Psi.$$

Proof of Lemma 4.1. Notice that in the equalities

$$\lim_{l \rightarrow 0} l^{-1} (T_{s+l}(f_i, \phi) - T_s(f_i, \phi)) = \lim_{l \rightarrow 0} l^{-1} \int_s^{s+l} T_m(Zf_i, \phi) dm = T_s(Zf_i, \phi)$$

we have the uniform convergence w.r.t. $i \in \mathbb{N}$ because there exists

$$\lim_{i \rightarrow \infty} Zf_i = h$$

and the family of the functions $\{T_l(\cdot, \phi) | l \in [0, s]\}$ is uniformly continuous. Therefore,

$$\begin{aligned} T_s(g, \phi) &= \lim_{i \rightarrow \infty} \lim_{l \rightarrow 0} l^{-1} (T_{s+l}(f_i, \phi) - T_s(f_i, \phi)) \\ &= \lim_{l \rightarrow 0} \lim_{i \rightarrow \infty} l^{-1} (T_{s+l}(f_i, \phi) - T_s(f_i, \phi)) \\ &= \lim_{l \rightarrow 0} l^{-1} (T_{s+l}(h, \phi) - T_s(h, \phi)) = (T_s(h, \phi))'_s. \end{aligned}$$

□

Lemma 4.2. Suppose that the assumptions of Proposition 3.1 are satisfied. If $f \in \mathcal{D}(\overline{F'(0)}) \cap \Phi$ and $\overline{F'(0)}f \in \Phi$, then

$$(T_s(f, \phi))'_s = T_s(\overline{F'(0)}f, \phi), \quad s \geq 0, \quad \phi \in \Psi.$$

Proof of Lemma 4.2. Since $f \in \mathcal{D}(\overline{F'(0)})$, we see that there exists a sequence $\{f_i\}_{i=1}^{\infty}$, $f_i \in \mathcal{D}(F'(0))$, such that $\lim_{i \rightarrow \infty} f_i = f$ and

$$\lim_{i \rightarrow \infty} F'(0)f_i = \overline{F'(0)}f.$$

Let Φ' be the minimal closed space such that $\Phi \subset \Phi'$ and $f_i, F'(0)f_i \in \Phi'$ for all $i \in \mathbb{N}$. Then, according to Proposition 3.1, we can choose a subsequence $\{m_k\}_{k=1}^{\infty}$ of the sequence $\{g_k\}_{k=1}^{\infty}$ such that we can define the family of the functions $T_s^1 : (\Phi', \Psi) \mapsto \mathbb{T}$, $s \geq 0$, by the equality

$$T_s^1(g, \phi) = \lim_{k \rightarrow \infty} (\{F(t/m_k)\}^{[m_k \frac{s}{t}]} g, \phi), \quad g \in \Phi', \quad \phi \in \Psi.$$

Thus, it follows from Lemma 4.1 (applied with parameters $h = f$, $g = \overline{F'(0)}f$) that

$$(T_s^1(f, \phi))'_s = T_s^1(\overline{F'(0)}f, \phi), \quad s \geq 0.$$

Since the restriction of T_s^1 to the set $\Phi \times \Psi$ is equal to T_s we have

$$(T_s(f, \phi))'_s = T_s(\overline{F'(0)}f, \phi), \quad s \geq 0, \quad \phi \in \Psi,$$

and the Lemma is proved. □

Proposition 4.1. Suppose that $F \in \mathcal{F}_{\mathbf{X}}$ and $t, l > 0$. Assume also that there exists a local solution $f : [0, l) \mapsto \mathcal{D}(\overline{F'(0)})$ of the system

$$(4.1) \quad f'(s) = \overline{F'(0)}f(s), \quad s \in [0, l),$$

$$(4.2) \quad f(0) = g \in \mathcal{D}(\overline{F'(0)}).$$

Then this solution is unique and

$$(4.3) \quad f(s) = w\text{-}\lim_{n \rightarrow \infty} \{F(t/n)\}^{[n\frac{s}{l}]} g, \quad s \in [0, l).$$

Proof of Proposition 4.1. Let $f(s)$, $s \in [0, l)$, be a local solution of system (4.1)-(4.2). Then for any $r \in \mathbb{Q} \cap [0, l)$ there exists a sequence $\{x_n^r\}_{n=1}^\infty$, $x_n^r \in \mathcal{D}(F'(0))$, such that

$$\lim_{n \rightarrow \infty} x_n^r = f(r)$$

and

$$\lim_{n \rightarrow \infty} \overline{F'(0)}x_n^r = \overline{F'(0)}f(r).$$

Let $\mathbf{B}$ be the minimal closed linear subspace of $\mathbf{X}$ such that $x_n^r \in \mathbf{B}$ for all $r \in \mathbb{Q} \cap [0, l)$, $n \in \mathbb{N}$. Hence, $f(s) \in \mathbf{B}$ for all $s \in [0, l)$ and, consequently, $\overline{F'(0)}f(s) \in \mathbf{B}$ for all $s \in [0, l)$. For any $\phi \in \mathbf{X}^*$ let us choose a subsequence $\{g_k\}_{k=1}^\infty$ of an arbitrary increasing sequence $\{n_k\}_{k=1}^\infty$, $n_k \in \mathbb{N}$, such that we can define the family of the functions $T_s : \Phi \times \Psi \mapsto \mathbb{T}$, $s \geq 0$, by equality (3.2) with $\Phi = \mathbf{B}$, $\Psi = \{\phi\}$. From Lemma 4.2 it follows that

$$(T_s(f(v), \phi))'_s = T_s(\overline{F'(0)}f(v), \phi), \quad s, v \in [0, l).$$

So, by Proposition 3.1,

$$\begin{aligned} (T_s(f(v-s), \phi))'_s &= \lim_{a \rightarrow 0} a^{-1} (T_{s+a}(f(v-a-s), \phi) - T_s(f(v-s), \phi)) \\ &= \lim_{a \rightarrow 0} a^{-1} (T_{s+a}((f(v-a-s) - f(v-s)), \phi)) \\ &+ \lim_{a \rightarrow 0} a^{-1} (T_{s+a}(f(v-s), \phi) - T_s(f(v-s), \phi)) \\ &= T_s(\overline{F'(0)}f(v-s), \phi) - T_s(\overline{F'(0)}f(v-s), \phi) = 0, \end{aligned}$$

for $v \in (0, l)$, $s \in (0, v)$. As a consequence, the function $T_s(f(v-s), \phi)$ is a constant w.r.t. s . Thus,

$$T_v(g, \phi) = (f(v), \phi), \quad v \in [0, l).$$

Now the existence of limit (4.3) follows from the arbitrariness of the sequence $\{n_k\}_{k=1}^\infty$ for any fixed $\phi \in \mathbf{X}^*$. Therefore, f is unique. $\square$

As a consequence of Proposition 4.1, we get

Theorem 4.1. Assume that $F \in \mathcal{F}_{\mathbf{X}}$ and $t, l > 0$. Assume also that a linear operator Z has the domain $\mathcal{D}(Z) \subset \mathcal{D}(F'(0))$ and

$$Zf = F'(0)f, \quad f \in \mathcal{D}(Z).$$

If there exists a local solution $f : [0, l) \mapsto \mathcal{D}(\overline{Z})$ of the system

$$f'(s) = \overline{Z}f(s), \quad s \in [0, l),$$

$$f(0) = g \in \mathcal{D}(\overline{Z}),$$

then the solution is unique and

$$f(s) = w\text{-}\lim_{n \rightarrow \infty} \{F(t/n)\}^{[n\frac{s}{t}]} g, \quad s \in [0, l].$$

Remark 4.1. The existence of the closure of Z in Theorem 4.1 follows from Theorem 3.1.

Corollary 4.1. Assume that the conditions of Theorem 4.1 are satisfied. Then

$$f'(s) = w\text{-}\lim_{n \rightarrow \infty} \{F(t/n)\}^{[n\frac{s}{t}]} \overline{Z}f(0), \quad s \in [0, l].$$

Proof of Corollary 4.1. Define the set $\mathbf{B}$ as in the proof of Proposition 4.1. For any $\phi \in \mathbf{X}^*$ choose a subsequence $\{g_k\}_{k=1}^\infty$ of an arbitrary increasing sequence $\{n_k\}_{k=1}^\infty$, $n_k \in \mathbb{N}$, such that we can define the family of the functions $T_s : \Phi \times \Psi \mapsto \mathbb{T}$, $s \geq 0$, by equality (3.2) with $\Phi = \mathbf{B}$, $\Psi = \{\phi\}$. From Lemma 4.2 and Theorem 4.1 it follows that

$$(f'(s), \phi) = \lim_{k \rightarrow \infty} (\{F(t/g_k)\}^{[g_k\frac{s}{t}]} \overline{Z}f(0), \phi), \quad s \in [0, l].$$

Since $\{n_k\}_{k=1}^\infty$ is arbitrary for any fixed $\phi \in \mathbf{X}^*$, the Corollary is proved. $\square$

5. CRITERIA FOR CLOSURE OF OPERATOR TO BE A GENERATOR OF C_0 -SEMIGROUP.

Lemma 5.1. Let $F \in \mathcal{F}_{\mathbf{X}}$. Assume that for some $t, l > 0$ and set $\Phi \subset \mathbf{X}$ there exists an increasing natural sequence $\{g_k\}_{k=1}^\infty$ such that there exists the limit

$$(5.1) \quad w\text{-}\lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k s]} g,$$

for any $g \in \Phi$ and any rational $s \in [0, l]$. Then limit (5.1) exists for any $g \in \overline{\text{span}\{\Phi\}}$, $s \in [0, l]$. Furthermore, the family of the operators $T_s : \overline{\text{span}\{\Phi\}} \mapsto \mathbf{X}$, $s \in [0, l]$, defined by

$$(5.2) \quad T_s g = w\text{-}\lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k\frac{s}{t}]} g,$$

satisfies the following conditions:

a) $T_s g$ is linear w.r.t. g and there exist $M \geq 1$ and $a \in \mathbb{R}$ such that

$$\|T_s g\| \leq M \exp(as) \|g\|, \quad s \in [0, l].$$

b) $T_s g$ is continuous w.r.t. $s \in [0, l]$ for all $g \in \overline{\text{span}\{\Phi\}}$.

c) If $f \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(F'(0))$ and $F'(0)f \in \overline{\text{span}\{\Phi\}}$, then there exists

$$(T_s f)'_s = T_s F'(0)f, \quad s \in [0, l].$$

d) If $\phi \in \mathcal{D}(F^{*'}(0))$, then

$$(T_m f, \phi) - (T_p f, \phi) = \int_p^m (T_s f, F^{*'}(0)\phi) ds, \quad m, p \in [0, l]$$

for any $f \in \overline{\text{span}\{\Phi\}}$.

e) If $\phi \in \mathcal{D}(F^{*'}(0))$, then $(T_s f, \phi)'_s = (T_s f, F^{*'}(0)\phi)$, $s \in [0, l]$, for any $f \in \overline{\text{span}\{\Phi\}}$.

f) If $f \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(F'(0))$, $F'(0)f \in \overline{\text{span}\{\Phi\}}$ and $\mathcal{D}(F^{*'}(0))$ is $*$ -dense in $\mathbf{X}^*$, then $T_s f \in \mathcal{D}((F^{*'}(0))^*)$, $s \in [0, l]$, and the following statements are valid:

1) $\mathcal{D}((F^{*'}(0))^*) \supset \mathcal{D}(F'(0))$, $(F^{*'}(0))^* f = F'(0)f$.

2) $(T_s f)'_s = (F^{*'}(0))^* T_s f$ for all $s \in [0, l]$.

Proof of Lemma 5.1. The existence of limit (5.1) for any $g \in \overline{\text{span}\{\Phi\}}$, $s \geq 0$ can be proved similarly to the first part of Proposition 3.1. Part (a) is trivial. Parts (b) – (e) easily follow from Proposition 3.1. Let us show part (f). We can notice that the following chain of the equalities can be deduced from parts (c) and (e)

$$((T_s f, g))'_s = (T_s F'(0)f, g) = (T_s f, F^{*'}(0)g), \quad g \in \mathcal{D}(F^{*'}(0)).$$

Thus, by *-density of $\mathcal{D}(F^{*'}(0))$ in $\mathbf{X}^*$, we get $T_s f \in \mathcal{D}((F^{*'}(0))^*)$ and $(F^{*'}(0))^* T_s f = T_s F'(0)f$ for each $s \geq 0$. From this it follows that $f \in \mathcal{D}((F^{*'}(0))^*)$ and $(F^{*'}(0))^* f = F'(0)f$. Hence, by part (c), we get that $(T_s f)'_s = (F^{*'}(0))^* T_s f$. $\square$

Theorem 5.1. Let $F \in \mathcal{F}_{\mathbf{X}}$. Assume that a linear operator Z has the domain $\mathcal{D}(Z) \subset \mathcal{D}(F'(0))$ and

$$Zf = F'(0)f, \quad f \in \mathcal{D}(Z).$$

Assume also that $\mathbf{A} \subset \mathcal{D}(Z)$ is a dense linear subset of $\mathbf{X}$ and there exists a fixed $l > 0$ such that there exists a local solution $f : [0, l) \mapsto \mathcal{D}(\bar{Z})$ of the system

$$(5.3) \quad f'(s) = \bar{Z}f(s), \quad s \in [0, l),$$

$$(5.4) \quad f(0) = f_0$$

for each $f_0 \in \mathbf{A}$. Then Z is closable and $\bar{Z}$ is a generator of C_0 -semigroup S . Furthermore, the following equality is satisfied:

$$(5.5) \quad S(t)f = \lim_{n \rightarrow \infty} F(t/n)^n f, \quad t \geq 0,$$

for all $f \in \mathbf{X}$.

Proof of Theorem 5.1. By Theorem 4.1, there exists the limit

$$(5.6) \quad w\text{-}\lim_{n \rightarrow \infty} F(t/n)^{[ns/t]} f, \quad s \in [0, l),$$

for each $f \in \mathbf{A}$. From the density of $\mathbf{A}$ in $\mathbf{X}$ and Lemma (5.1) we easily infer that there exists limit (5.6) for each $f \in \mathbf{X}$. Put $G(s)f = w\text{-}\lim_{n \rightarrow \infty} F(t/n)^{[ns/t]} f$, $s \in [0, l)$, $f \in \mathbf{X}$. From the uniqueness of the local solution of system (5.3)–(5.4) we infer that

$$(5.7) \quad G(s_1)G(s_2)f = G(s_1 + s_2)f, \quad s_1, s_2, s_1 + s_2 \in [0, l),$$

for each $f \in \mathbf{A}$. From this and part (a) of Lemma (5.1) it follows that equality (5.7) is valid for each $f \in \mathbf{X}$. Define the function $S : [0, \infty) \mapsto \mathcal{L}(\mathbf{X})$ by the equality

$$S(s) = \{G(l/2)\}^{[2s/l]} G(s - [2s/l]l/2), \quad s \geq 0.$$

Then, from parts (a), (b) of Lemma (5.1) and the definition of S we easily infer that the function S is C_0 -semigroup. Furthermore, from part (c) of Lemma (5.1) it follows that $\mathcal{D}(Z) \subset \mathcal{D}(F'(0)) \subset \mathcal{D}(S'(0))$ and, by the closedness of $S'(0)$, $\bar{Z}f = S'(0)f$ for each $f \in \mathcal{D}(\bar{Z})$. Let us show that $\mathcal{D}(\bar{Z}) = \mathcal{D}(S'(0))$. Fix $g \in \mathbf{A}$. Then

$$\sum_{k=1}^n \frac{s}{n} S(k \frac{s}{n}) g \in \mathbf{A}$$

and, by properties of C_0 -semigroups, there exist the limits

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{s}{n} S(k \frac{s}{n}) g &= \int_0^s S(t) g dt, \\ \lim_{n \rightarrow \infty} \bar{Z} \sum_{k=1}^n \frac{s}{n} S(k \frac{s}{n}) g &= \int_0^s S(t) \bar{Z} g dt = S(s)g - g, \quad s \in [0, l/2]. \end{aligned}$$

Thus, by the closedness of $\bar{Z}$, we get that $\int_0^s S(t) g dt \in \mathcal{D}(\bar{Z})$ and

$$\bar{Z} \int_0^s S(t) g dt = S(s)g - g.$$

Let $f \in \mathbf{X}$. Choose a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \mathbf{A}$, such that $\lim_{n \rightarrow \infty} f_n = f$. Then, by properties of C_0 -semigroups, there exist the limits

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_0^s S(t) f_n dt &= \int_0^s S(t) f dt, \\ \lim_{n \rightarrow \infty} \bar{Z} \int_0^s S(t) f_n dt &= \lim_{n \rightarrow \infty} S(s) f_n - f_n = S(s)f - f, \quad s \in [0, l/2]. \end{aligned}$$

Then, by the closedness of $\bar{Z}$,

$$\int_0^s S(t) f dt \in \mathcal{D}(\bar{Z})$$

and

$$\bar{Z} \int_0^s S(t) f dt = S(s)f - f$$

for $s \in [0, l/2]$. Notice that for $f \in \mathcal{D}(S'(0))$ there exist the limits

$$\begin{aligned} \lim_{s \rightarrow 0} s^{-1} \int_0^s S(t) f dt &= f, \\ \lim_{s \rightarrow 0} s^{-1} \bar{Z} \int_0^s S(t) f dt &= \lim_{s \rightarrow 0} s^{-1} (S(s)f - f) = S'(0)f. \end{aligned}$$

So, by the closedness of $\bar{Z}$, $f \in \mathcal{D}(\bar{Z})$. Thus, $\bar{Z}$ is a generator of C_0 -semigroup S and, by Chernoff's theorem, we get equality (5.5). $\square$

Theorem 5.2. Let Z be a densely defined linear operator in $\mathbf{X}$ and $t > 0$. Then Z is closable and its closure is a generator of C_0 -semigroup if and only if there exists a function $F \in \mathcal{F}_{\mathbf{X}}$ such that:

- i) $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$.
- ii) $\mathcal{D}(F^{*'}(0))$ is $*$ -dense in $\mathbf{X}^*$.
- iii) There exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(Z)$ such that for any $f \in \mathbf{A}$, $s \geq 0$ there exists a subsequence $\{f_n^s\}_{n=1}^\infty$, $f_n^s \in \mathcal{D}(Z)$, that satisfies the following conditions:
 - a) $\lim_{n \rightarrow \infty} \|F^{[ns]}(\frac{t}{n})f - f_n^s\| = 0$.
 - b) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{f_n^s\}_{n \in \mathbb{N}}$.
 - c) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{Zf_n^s\}_{n \in \mathbb{N}}$.

Furthermore, if conditions (i)-(iii) are satisfied, then $\exp(s\bar{Z})f = \lim_{n \rightarrow \infty} F(s/n)^n f$, $s > 0$, for all $f \in X$.

Proof of Theorem 5.2. If $\bar{Z}$ is a generator of C_0 -semigroup and $F(s) = \exp(s\bar{Z})$, $s \geq 0$, then it is easy to check that conditions (i)-(iii) are satisfied. Indeed, it follows from Statement 2.1 that conditions (i), (ii) are satisfied. Put $\mathbf{A} = \mathcal{D}(Z)$. We can notice that if $f \in \mathbf{A}$, then $\exp(s\bar{Z})f \in \mathcal{D}(\bar{Z})$. From this we infer condition (iii). Now let us show that conditions (i)-(iii) imply that $\bar{Z}$ is a generator of C_0 -semigroup. Let $\{s_k\}_{k=1}^\infty$ be a sequence of all nonnegative rational numbers. Fix $f \in \mathbf{A}$ and an increasing sequence $\{n_i\}_{i=1}^\infty$ of natural numbers. For each $i \in \mathbb{N}$ let the sequence $\{f_n^{s_i}\}_{n=1}^\infty$ satisfy condition (iii) for s_i and f . Choose a subsequence $\{n_i^1\}_{i=1}^\infty$ of the sequence $\{n_i\}_{i=1}^\infty$ such that there exist the limits $w\text{-}\lim_{i \rightarrow \infty} f_{n_i^1}^{s_1}$ and $w\text{-}\lim_{i \rightarrow \infty} Z f_{n_i^1}^{s_1}$. Then, it follows from part (a) of condition (iii) that there exists the limit $w\text{-}\lim_{i \rightarrow \infty} F^{[n_i^1 s_1]}(t/n_i^1)f$. Similarly, choose a subsequence $\{n_i^2\}$ of the sequence $\{n_i^1\}$ such that there exist the limits $w\text{-}\lim_{i \rightarrow \infty} f_{n_i^2}^{s_2}$, $w\text{-}\lim_{i \rightarrow \infty} Z f_{n_i^2}^{s_2}$, $w\text{-}\lim_{i \rightarrow \infty} F^{[n_i^2 s_2]}(t/n_i^2)f$. In the same way, for any natural $k \geq 3$, we choose a subsequence $\{n_i^k\}_{i=1}^\infty$ of the sequence $\{n_i^{k-1}\}_{i=1}^\infty$ such that there exist the limits $w\text{-}\lim_{i \rightarrow \infty} f_{n_i^k}^{s_k}$, $w\text{-}\lim_{i \rightarrow \infty} Z f_{n_i^k}^{s_k}$, $w\text{-}\lim_{i \rightarrow \infty} F^{[n_i^k s_k]}(t/n_i^k)f$. Then we can consider the diagonal sequence $\{n_i^i\}_{i=1}^\infty$ and deduce that there exist the limits $w\text{-}\lim_{i \rightarrow \infty} f_{n_i^i}^{s_i}$, $w\text{-}\lim_{i \rightarrow \infty} Z f_{n_i^i}^{s_i}$, $w\text{-}\lim_{i \rightarrow \infty} F^{[n_i^i s_i]}(t/n_i^i)f$ for any $k \in \mathbb{N}$. Choose now a sequence $\{g_l\}_{l=1}^\infty$, $g_l \in \mathbf{A}$, such that $\lim_{k \rightarrow \infty} g_k = F'(0)f$. Choose a subsequence d_k of the sequence $\{n_k^k\}_{k=1}^\infty$ such that there exists the limit $w\text{-}\lim_{l \rightarrow \infty} F^{[d_l s_k]}(t/d_l)g_m$ for any $k, m \in \mathbb{N}$. Let

$$\mathbf{B} = \overline{\text{span} \{ \{f\} \cup \{g_k | k \in \mathbb{N}\} \}}.$$

Then it follows from Lemma 5.1 that there exists the limit $w\text{-}\lim_{i \rightarrow \infty} F^{[d_i s]}(t/d_i)g$ for any $s \geq 0$, $g \in \mathbf{B}$. Define

$$T_s g = w\text{-}\lim_{i \rightarrow \infty} F^{[d_i s]}(t/d_i)g, \quad g \in \mathbf{B}.$$

From the coincidence of the weak closure of Z with the closure in the strong topology it follows that $T_{s_k} f \in \mathcal{D}(\bar{Z})$ for any $k \in \mathbb{N}$ because there exist the limits $w\text{-}\lim_{i \rightarrow \infty} f_{d_i}^{s_k}$, $w\text{-}\lim_{i \rightarrow \infty} Z f_{d_i}^{s_k}$. From part (f) of Lemma 5.1 and the closedness of the operator $(F^{*'}(0))^*$ we infer that

$$(T_s f)'_s = \bar{Z} T_s f$$

for any rational $s \geq 0$. Further, we can choose for arbitrary $m > 0$ a sequence $\{m_i\}$, $m_i \in \mathbb{Q} \cap [0, \infty)$, such that $\lim_{i \rightarrow \infty} m_i = m$. Then

$$\lim_{i \rightarrow \infty} T_{m_i} f = T_m f$$

and

$$\lim_{i \rightarrow \infty} \bar{Z} T_{m_i} f = \lim_{i \rightarrow \infty} (T_s f)'_{m_i} f = \lim_{i \rightarrow \infty} T_{m_i} Z f = T_m Z f.$$

Therefore, by the closability of Z , $T_m f \in \mathcal{D}(\bar{Z})$ and

$$(T_s f)'_s = \bar{Z} T_s f$$

for all $s \geq 0$. Consequently, $T_s f$ is a solution of the equation

$$f'(s) = \bar{Z}f(s)$$

with the initial condition $f(0) = f$. Thus, Theorem 5.1 implies that $\bar{Z}$ is the generator of C_0 -semigroup S such that

$$S(s)f = \lim_{n \rightarrow \infty} F(s/n)^n f, \quad s \geq 0,$$

for each $f \in X$. □

Theorem 5.3. Let Z be a densely defined linear operator in $\mathbf{X}$ and $t > 0$. Assume also that there exists a function $F \in \mathcal{F}_X$ such that:

- i) $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$.
- ii) $\mathcal{D}(F^{*'}(0))$ *-dense in $\mathbf{X}^*$.

Then operator Z is closable and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(Z)$ such that for all $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n^s\}_{n=1}^\infty$, $f_n^s \in \mathcal{D}(Z)$, satisfying the following conditions:

- a) $\lim_{n \rightarrow \infty} \|F^{[ns]}(\frac{t}{n})f - f_n^s\| = 0$.
- b) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{Zf_n^s\}_{n \in \mathbb{N}}$.
- c) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{f_n^s\}_{n \in \mathbb{N}}$.

Proof of Theorem 5.3. ($\Rightarrow$) If operator Z is closable and its closure is a generator of C_0 -semigroup, then, by Chernoff's Theorem, it is enough to put $\mathbf{A} = \mathcal{D}(Z)$ and choose sequences $\{f_n^s\}_{n=0}^\infty$, $f_n^s \in \mathbf{A}$, $s \geq 0$, such that there exist the limits $\lim_{n \rightarrow \infty} f_n^s = \exp(s\bar{Z})f$,

$$\lim_{n \rightarrow \infty} Zf_n^s = \bar{Z} \exp(s\bar{Z})f.$$

($\Leftarrow$) It follows from Theorem 5.2. □

Corollary 5.1. Let $t > 0$. Assume that C and D are generators of C_0 -semigroups $\exp(sC)$ and $\exp(sD)$ in $\mathbf{X}$. Assume also that there exists a set $\mathbf{B} \subset \mathcal{D}((\exp(sC)^*)'_{s=0}) \cap \mathcal{D}((\exp(sD)^*)'_{s=0})$ and the following conditions are satisfied:

- i) $\mathcal{D}(C) \cap \mathcal{D}(D)$ is dense in $\mathbf{X}$.
- ii) $\mathbf{B}$ is *-dense in $\mathbf{X}^*$.
- iii) There exist $a \in \mathbb{R}$ and $M \geq 1$ such that

$$\|\{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^m\| \leq M \exp(as\frac{m}{n})$$

for all $n, m \in \mathbb{N}$, $s > 0$.

- iv) Function $g(s, x) = \exp(sD^*)x$, $s \geq 0$, $x \in \mathbf{X}$, is continuous at $s = 0$ for $x \in (\exp(sC)^*)'_{s=0}(\mathbf{B})$.

Then the sum of C and D is a closable operator and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(C) \cap \mathcal{D}(D)$ such that for all $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n^s\}_{n=1}^\infty$, $f_n^s \in \mathcal{D}(C) \cap \mathcal{D}(D)$, satisfying the following conditions:

- a) $\lim_{n \rightarrow \infty} \|(\exp(\frac{t}{n}C) \exp(\frac{t}{n}D))^{[ns]}f - f_n^s\| = 0$.
- b) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{f_n^s\}_{n=1}^\infty$.

c) It is possible to choose a weakly convergent subsubsequence for any subsequence of the sequence $\{(C + D)f_n^s\}_{n=1}^\infty$.

Furthermore, if conditions (a)-(c) are satisfied, then

$$\exp(s(\overline{C + D}))f = \lim_{n \rightarrow \infty} \{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^n f, \quad s \geq 0,$$

for all $f \in X$.

Proof of Corollary 5.1. Let $F(s) = \exp(sC) \exp(sD)$. Since

$$\begin{aligned} \lim_{s \rightarrow 0} s^{-1}(F(s)f - F(0)f) &= \lim_{s \rightarrow 0} s^{-1}(\exp(sC) \exp(sD)f - \exp(sC)f) \\ &+ \lim_{s \rightarrow 0} s^{-1}(\exp(sC)f - f) = Cf + Df, \quad f \in \mathcal{D}(C) \cap \mathcal{D}(D), \end{aligned}$$

and

$$\begin{aligned} \lim_{s \rightarrow 0} s^{-1}(F^*(s) - F(0))f &= \lim_{s \rightarrow 0} s^{-1}\{(\exp(sD))^*(\exp(sC))^* - (\exp(sD))^*\}f \\ &+ \lim_{s \rightarrow 0} s^{-1}\{(\exp(sD))^*f - f\} = \lim_{l \rightarrow 0} (\exp(lD))^*(\exp(sC)^*)'|_{s=0}f \\ &+ (\exp(sD)^*)'|_{s=0}f = (\exp(sC)^*)'|_{s=0}f + (\exp(sD)^*)'|_{s=0}f, \quad f \in B, \end{aligned}$$

we infer that $\mathcal{D}(F'(0))$ is dense in X and $\mathcal{D}(F^*(0))$ is $*$ -dense in X^* . Therefore, we can apply Theorem 5.3 with $Z = C + D$. $\square$

Lemma 5.2. Let $F \in \mathcal{F}_X$. Assume that for some $t > 0$ and set $\Phi \subset X$ there exists an increasing natural sequence $\{g_k\}_{k=1}^\infty$ such that there exists the limit

$$(5.8) \quad \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k s]} g$$

for any $g \in \Phi$ and any nonnegative rational s . Then limit (5.8) exists for any $g \in \overline{\text{span}\{\Phi\}}$, $s \geq 0$. Furthermore, the family of the operators $T_s : \overline{\text{span}\{\Phi\}} \mapsto X$, $s \geq 0$, defined by

$$T_s g = \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k \frac{s}{t}]} g,$$

satisfies the following conditions:

a) If $T_s f \in \overline{\text{span}\{\Phi\}}$ for some $s \geq 0$, then

$$T_{s+l} f = T_l T_s f, \quad l > 0.$$

b) Let $f \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(\overline{F'(0)})$ and $\overline{F'(0)}f \in \overline{\text{span}\{\Phi\}}$. Assume also that there exists a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(F'(0))$, such that $\lim_{n \rightarrow \infty} f_n = f$ and $\lim_{n \rightarrow \infty} F'(0)f_n = \overline{F'(0)}f$. Then there exists

$$(T_s f)'_s = T_s \overline{F'(0)}f, \quad s \geq 0.$$

c) Let $T_s f \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(\overline{F'(0)})$, $f \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(F'(0))$ and $\overline{F'(0)}T_s f, F'(0)f \in \overline{\text{span}\{\Phi\}}$ for some $s \geq 0$. Assume also that there exists a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \overline{\text{span}\{\Phi\}} \cap \mathcal{D}(F'(0))$, such that $\lim_{n \rightarrow \infty} f_n = T_s f$ and $\lim_{n \rightarrow \infty} F'(0)f_n = \overline{F'(0)}T_s f$. Then there exists

$$(T_s f)'_s = \overline{F'(0)}T_s f, \quad s \geq 0.$$

Proof of Lemma 5.2. The existence of limit (5.8) for any $g \in \overline{\text{span}\{\Phi\}}$, $s \geq 0$ can be proved similarly to the first part of Proposition 3.1.

(a) Part (a) follows from the chain of the equalities:

$$\begin{aligned} T_l T_s f &= \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k l]} T_s f = \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k l]} (T_s - \{F(t/g_k)\}^{[g_k s]}) f \\ &\quad + \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k l]} \{F(t/g_k)\}^{[g_k s]} f = \lim_{k \rightarrow \infty} \{F(t/g_k)\}^{[g_k l] + [g_k s]} f = T_{l+s} f \end{aligned}$$

(b) By part (c) of Lemma 5.1, we have

$$T_s f - T_r f = \lim_{n \rightarrow \infty} T_s f_n - T_r f_n = \lim_{n \rightarrow \infty} \int_r^s T_l F'(0) f_n dl = \int_r^s T_l \overline{F'(0)} f dl$$

for any $r, s \geq 0$. From this we easily infer part (b).

(c) From part (c) of Lemma 5.1 and parts (a), (b) of Lemma 5.2 it follows that

$$(T_s f)'_s = T_s F'(0) f = \lim_{l \rightarrow 0} l^{-1} (T_{s+l} f - T_s f) = \lim_{l \rightarrow 0} l^{-1} (T_l - I) T_s f = \overline{F'(0)} T_s f.$$

□

The following two results are respectively extensions of Corollaries 4, 5 from [4].

Theorem 5.4. Let Z be a densely defined linear operator in $\mathbf{X}$ and $t > 0$. Assume also that there exists a function $F \in \mathcal{F}_{\mathbf{X}}$ such that $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$. Then operator Z is closable and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(Z)$ such that for all $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n^s\}_{n=1}^\infty$, $f_n^s \in \mathcal{D}(Z)$, satisfying the following conditions:

a) $\lim_{n \rightarrow \infty} \|F^{[ns]}(\frac{t}{n})f - f_n^s\| = 0$.

b) The sets $\{Zf_n^s\}_{n=1}^\infty$ and $\{f_n^s\}_{n=1}^\infty$ are precompact.

Proof of Theorem 5.4. ($\Rightarrow$) If operator Z is closable and its closure is a generator of C_0 -semigroup, then, by Chernoff's Theorem, it is enough to put $\mathbf{A} = \mathcal{D}(Z)$ and choose sequences $\{f_n^s\}_{n=0}^\infty$, $f_n^s \in \mathbf{A}$, $s \geq 0$, such that there exist the limits $\lim_{n \rightarrow \infty} f_n^s = \exp(s\bar{Z})f$,

$$\lim_{n \rightarrow \infty} Zf_n^s = \bar{Z} \exp(s\bar{Z})f.$$

($\Leftarrow$) From condition (a) it follows that $\{F^{[ns]}(\frac{t}{n})f\}_{n=1}^\infty$ is precompact for each $f \in \mathbf{A}$ and $s > 0$. Then, by the density $\mathbf{A}$ in $\mathbf{X}$, $\{F^{[ns]}(\frac{t}{n})f\}_{n=1}^\infty$ is precompact for each $f \in \mathbf{X}$ and $s > 0$. Fix $f \in \mathbf{A}$. As in the proof of Theorem 5.2, we can choose an increasing sequence $\{g_n\}_{n=1}^\infty$ such that there exist the limits $\lim_{n \rightarrow \infty} F^{[g_n s]}(\frac{t}{g_n})f$, $\lim_{n \rightarrow \infty} Zf_{g_n}^s$, $\lim_{n \rightarrow \infty} F^{[g_n l]}(\frac{t}{g_n})f_k^s$ and $\lim_{n \rightarrow \infty} F^{[g_n l]}(\frac{t}{g_n})Zf_k^s$ for each $s, l \in [0, \infty) \cap \mathbb{Q}$ and $k \in \mathbb{N}$. Let

$$\mathbf{B} = \overline{\text{span} \{ \{f\} \cup \{Zf_k^s | k \in \mathbb{N}, s \in [0, \infty) \cap \mathbb{Q}\} \cup \{f_k^s | k \in \mathbb{N}, s \in [0, \infty) \cap \mathbb{Q}\} \}}.$$

Then it follows from Lemma 5.2 that there exists the limit $\lim_{i \rightarrow \infty} F^{[d_i s]}(t/d_i)g$ for any $s \geq 0$, $g \in \mathbf{B}$. Define

$$T_s g = \lim_{i \rightarrow \infty} F^{[d_i s]}(t/d_i)g, \quad g \in \mathbf{B}.$$

By condition (a), there exist the limits $\lim_{i \rightarrow \infty} f_{g_n}^s = T_s f$ and $\lim_{i \rightarrow \infty} Zf_{g_n}^s$ for each $s \in [0, \infty) \cap \mathbb{Q}$. So, by the closability of Z , $T_s f \in \mathcal{D}(\bar{Z}) \cap \mathbf{B}$ and

$$\lim_{i \rightarrow \infty} Zf_{g_n}^s = \overline{F'(0)} T_s f \in \mathbf{B}$$

for any $s \in [0, \infty) \cap \mathbb{Q}$. Thus, by part (c) of Lemma 5.2, we get

$$(T_s f)'_s = \bar{Z} T_s f$$

for any rational $s \geq 0$. Choose for arbitrary $m > 0$ a sequence $\{m_i\}$, $m_i \in \mathbb{Q} \cap [0, \infty)$, such that $\lim_{i \rightarrow \infty} m_i = m$. Then

$$\lim_{i \rightarrow \infty} T_{m_i} f = T_m f$$

and

$$\lim_{i \rightarrow \infty} \bar{Z} T_{m_i} f = \lim_{i \rightarrow \infty} (T_s f)'_{m_i} f = \lim_{i \rightarrow \infty} T_{m_i} Z f = T_m Z f.$$

Therefore, by the closability of Z , $T_m f \in \mathcal{D}(\bar{Z})$ and

$$(T_s f)'_s = \bar{Z} T_s f$$

for all $s \geq 0$. Consequently, $T_s f$ is a solution of the equation

$$f'(s) = \bar{Z} f(s)$$

with the initial condition $f(0) = f$. Thus, Theorem 5.1 implies that $\bar{Z}$ is a generator of C_0 -semigroup and, by Chernoff's theorem,

$$\exp(s\bar{Z})f = \lim_{n \rightarrow \infty} F(s/n)^n f, \quad f \in X.$$

□

From Theorem 5.4 we get the following result.

Corollary 5.2. Let $t > 0$. Assume that C and D are generators of C_0 -semigroups $\exp(sC)$ and $\exp(sD)$. Assume also that the following conditions are satisfied:

- i) $\mathcal{D}(C) \cap \mathcal{D}(D)$ is dense in $\mathbf{X}$.
- ii) There exists $a \in \mathbb{R}$ and $M \geq 1$ such that

$$\|\{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^m\| \leq M \exp(as\frac{m}{n})$$

for all $n, m \in \mathbb{N}$, $s \in \mathbb{R}$.

Then the sum of C and D is a closable operator and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(C) \cap \mathcal{D}(D)$ such that for all $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n^s\}_{n=1}^\infty$, $f_n^s \in \mathcal{D}(C) \cap \mathcal{D}(D)$, satisfying the following conditions:

- a) $\lim_{n \rightarrow \infty} \|(\exp(\frac{t}{n}C) \exp(\frac{t}{n}D))^{[ns]} f - f_n^s\| = 0$.
- b) The sets $\{f_n^s\}_{n=1}^\infty$ and $\{(C+D)f_n^s\}_{n=1}^\infty$ are precompact.

Furthermore, if conditions (a)-(b) are satisfied, then

$$\exp(s(\overline{C+D}))f = \lim_{n \rightarrow \infty} \{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^n f, \quad s > 0,$$

for all $f \in X$.

If $\mathbf{X}$ is a reflexive separable Banach space or a Hilbert space, then the formulations of Theorems 5.2, 5.3 and Corollary 5.1 can be considerably simplified and, by Statement 2.2, we get the following Corollaries of the results mentioned above:

Corollary 5.3. Let $Z : \mathcal{B} \supset \mathcal{D}(Z) \rightarrow \mathcal{B}$ be a linear operator, $\mathcal{D}(Z)$ be dense in $\mathcal{B}$ and $t > 0$. Operator Z is closable and its closure is a generator of C_0 -semigroup iff there exists $F : [0, \infty) \mapsto \mathcal{L}(\mathcal{B})$ such that:

- i) $F(0) = I$ and there exists $a \in \mathbb{R}$, $M \geq 1$ such that $\|F^m(\frac{s}{n})\| \leq M \exp(as\frac{m}{n})$ for all $n, m \in \mathbb{N}$ and $s \geq 0$.
- ii) $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$.
- iii) $\mathcal{D}(F^{*'}(0))$ is $*$ -dense in $\mathcal{B}^*$.
- iv) There exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(Z)$ such that for any $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \mathcal{D}(Z)$ satisfying the following identity

$$\lim_{n \rightarrow \infty} \|F^{[ns]}(\frac{t}{n})f - f_n\| = 0$$

and, furthermore, $\{Zf_n\}_{n=1}^\infty$ is a bounded sequence.

Furthermore, if conditions (i)-(iv) are satisfied, then $\exp(s\overline{Z})f = \lim_{n \rightarrow \infty} F(s/n)^n f$, $s > 0$, for all $f \in \mathcal{B}$.

Corollary 5.4. Let $Z : \mathcal{B} \supset \mathcal{D}(Z) \rightarrow \mathcal{B}$ be a linear operator, $\mathcal{D}(Z)$ be dense in $\mathcal{B}$ and $t > 0$. Assume that there exists $F : [0, \infty) \mapsto \mathcal{L}(\mathcal{B})$ such that the following conditions are satisfied:

- i) $F(0) = I$ and there exists $a \in \mathbb{R}$, $M \geq 1$ such that

$$\|F^m(\frac{s}{n})\| \leq M \exp(as\frac{m}{n})$$

for all $n, m \in \mathbb{N}$ and $s \geq 0$.

- ii) $\mathcal{D}(F'(0)) \supset \mathcal{D}(Z)$ and $F'(0)f = Zf$, $f \in \mathcal{D}(Z)$.
- iii) $\mathcal{D}(F^{*'}(0))$ is $*$ -dense in $\mathcal{B}^*$.

Then operator Z is a closable operator and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(Z)$ such that for any $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \mathcal{D}(Z)$, satisfying the following identity $\lim_{n \rightarrow \infty} \|F^{[ns]}(\frac{t}{n})f - f_n\| = 0$ and, furthermore, $\{F'(0)f_n\}_{n=1}^\infty$ is a bounded sequence.

Corollary 5.5. Let $t > 0$. Suppose that C and D are generators of C_0 -semigroups $\exp(sC)$ and $\exp(sD)$ in $\mathcal{B}$. Assume also that the following conditions are satisfied:

- i) $\mathcal{D}(C) \cap \mathcal{D}(D)$ is dense in $\mathcal{B}$.
- ii) $\mathcal{D}(C^*) \cap \mathcal{D}(D^*)$ is dense in $\mathcal{B}^*$.
- iii) There exists $a \in \mathbb{R}$, $M \geq 1$ such that the following inequality holds

$$\|\{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^m\| \leq M \exp(as\frac{m}{n})$$

for all $n, m \in \mathbb{N}$, $s > 0$.

Then the sum of operators C and D is a closable operator and its closure is a generator of C_0 -semigroup iff there exists a dense linear subspace $\mathbf{A} \subseteq \mathcal{D}(C) \cap \mathcal{D}(D)$ such that for any $f \in \mathbf{A}$, $s \geq 0$ there exists a sequence $\{f_n\}_{n=1}^\infty$, $f_n \in \mathcal{D}(C) \cap \mathcal{D}(D)$, satisfying the following conditions:

- a) $\lim_{n \rightarrow \infty} \|(\exp(\frac{t}{n}C) \exp(\frac{t}{n}D))^{[ns]}f - f_n\| = 0$.
- b) $\{(C + D)f_n\}_{n=1}^\infty$ is a bounded sequence.

Furthermore, if conditions (a)-(b) are satisfied, then

$$\exp(s(\overline{C + D}))f = \lim_{n \rightarrow \infty} \{\exp(\frac{s}{n}C) \exp(\frac{s}{n}D)\}^n f, \quad s > 0,$$

for all $f \in \mathcal{B}$.

Remark 5.1. If operators iC and iD are self-adjoint, then conditions (ii), (iii) in Corollary 5.5 can be omitted.

REFERENCES

- [1] P. R. Chernoff, Note on product formulas for operator semigroups, J. Funct. Anal. Vol. 84, 238–242 (1968).
- [2] E. B. Davies, One-Parameter Semigroups, St. John's College, Oxford, England (1980).
- [3] K. J. Engel and R. Nagel, One-Parameter Semigroups for Linear Evolution Equations, Graduate Texts in Math., vol. 194, Springer-Verlag, 2000.
- [4] A. Y. Neklyudov, Inversion of Chernoff's theorem, accepted for publication in Mathematical Notes.
- [5] O. G. Smolyanov, A. G. Tokarev, A. Truman, Hamiltonian Feynman path integral via the Chernoff formula, J. Math. Phys., Vol. 43, No. 10, October 2002.
- [6] O. G. Smolyanov, H. von Weizsacker, O. Wittich, Brownian Motion on a Manifold as Limit of Step-wise Conditioned Standard Brownian Motions, Canadian Math. Society Conference Proceedings, 589-602, 29 (2000).
- [7] O. G. Smolyanov, H. von Weizsacker, O. Wittich, Diffusion on a compact Riemannian manifold and surface measures, (Russian) Dokl. Akad. Nauk 371 (2000), no. 4, 442–447.
- [8] O. G. Smolyanov, E. T. Shavgulidze, Path integrals, (Russian) Moskov. Gos. Univ., Moscow, 1990. 152 pp. ISBN: 5-211-00944-4